


TAUP 2013  
Asilomar, California 06-13 September 2013


# SELF-INDUCED FLAVOR EVOLUTION OF SUPERNOVA NEUTRINOS WITHOUT AXIAL SYMMETRY

(Based on 1308.1402; 1308.5255)

(Alessandro MIRIZZI, Hamburg U.)



# SNAP-SHOTS OF SN DENSITY PROFILES



- Matter bkg potential

$$\lambda = \sqrt{2} G_F N_e \sim R^{-3}$$

- $\nu$ - $\nu$  interaction

$$\mu = \sqrt{2} G_F n_\nu \sim R^{-2}$$

- Vacuum oscillation frequencies

$$\omega = \frac{\Delta m^2}{2E}$$

When  $\mu \gg \lambda$ , SN  $\nu$  oscillations dominated by  $\nu$ - $\nu$  interactions

Collective flavor transitions at low-radii [O ( $10^2 - 10^3$  km)]

Two seminal papers in 2006 triggered a torrent of activities  
 Duan, Fuller, Qian, astro-ph/0511275, Duan et al. astro-ph/0606616

[see Duan et al, arXiv:1001.2799 for a review]

# DENSITY MATRIX FOR THE NEUTRINO ENSEMBLE

Diagonal elements  
related to flavor  
content

$$\rho_{\alpha\alpha} = \frac{F_{\nu_\alpha}(E, r)}{F(E, r)}$$

$$\rho = \begin{pmatrix} \rho_{ee} & \rho_{e\mu} & \rho_{e\tau} \\ \rho_{e\mu}^* & \rho_{\mu\mu} & \rho_{\mu\tau} \\ \rho_{e\tau} & \rho_{\mu\tau}^* & \rho_{\tau\tau} \end{pmatrix}$$

Off-diagonal elements  
responsible for flavor  
conversions

- In 2 $\nu$  scenario ( $\Delta m^2_{\text{atm}}, \theta_{13}$ ). Decompose density matrix over Pauli matrices to get the "**polarization**" (Bloch) vector  $\mathbf{P}$ . Survival probability  $P_{ee} = 1/2(1 + \mathbf{P}_z)$ .  
 $\mathbf{P}_z = -1 \rightarrow P_{ee} = 0$  ;  $\mathbf{P}_z = 0 \rightarrow P_{ee} = 1/2$  (flavor decoherence)

- The EOMs for the time evolution in a homogeneous medium are the Liouville equations (e.g. Early Universe)

$$i\partial_t \rho_\varpi = [H_\varpi, \rho_\varpi]$$

# MULTI-ANGLE (M.A.) EOMs FOR SN NEUTRINOS


Evolution in space for  $\nu$ 's streaming from a SN core in quasi-stationary situation

$$i \vec{v}_p \cdot \vec{\nabla}_x \rho_{p,x} = [H(\omega, \lambda, \rho_{p',x}), \rho_{p,x}]$$

Liouville operator for free streaming  $\nu$


## MULTI-ANGLE $\nu$ - $\nu$ HAMILTONIAN

$$H_{\nu\nu} = \sqrt{2}G_F \int d\vec{q} (1 - \vec{v}_p \cdot \vec{v}_q) (\rho_{q,x} - \bar{\rho}_{q,x})$$



# BULB MODEL

[see, e.g., Duan et al., astro-ph/0606616] → First large-scale multi-angle simulations



- Neutrinos are emitted uniformly and (half)-isotropically from the surface of a sphere ( $\nu$ -sphere), like in a blackbody.
- Physical conditions depend only on the distance  $r$  from the center of the star (**azimuthal symmetry**)
- Only **multi-zenith-angle (MZA) effects** in terms of  $u = \sin^2 \theta_R$
- Project evolution along radial direction (ODE problem)  $\vec{\nabla}_p \cdot \vec{\nabla}_x \rightarrow v_r d_r$

# SINGLE-ZENITH-ANGLE (SZA) APPROXIMATION


Introduce an average of the MZA factor ( $1 - \cos\theta_{pq}$ ).

**SZA  $\nu$ - $\bar{\nu}$  Hamiltonian**

$$H_{\nu\bar{\nu}} = \mu(r)(\rho - \bar{\rho})$$

$$\mu(r) = \sqrt{2}G_F n_\nu \langle 1 - \cos\theta \rangle$$

*[Hannestad et al., astro-ph/0608695]*



**Toy model:**

Monochromatic  $\nu$  beam.

Only  $\nu_e$  and  $\bar{\nu}_e$ , with an excess of  $\nu_e$


- No effect in NH
- Complete self-induced conversions in IH. **Bimodal instability**

# MULTI-ZENITH-ANGLE DECOHERENCE

$$H_{\nu\nu} = \sqrt{2}G_F \int d\vec{q}^r \left(1 - \cos \theta_{pq}\right) \left(\rho_{q,x} - \bar{\rho}_{q,x}\right)$$

Flux term

Symmetric  $\nu_e \bar{\nu}_e$  systems decoheres in both hierarchies [Raffelt & Sigl, hep-ph/0701182]



Flux term does not vanish in a non-isotropic medium, like  $\nu$  streaming off a SN


Is the MZA decoherence relevant for SN neutrinos?

# MZA EFFECTS FOR SN NEUTRINOS

$$\varepsilon = \frac{F(\nu_e) - F(\bar{\nu}_e)}{F(\bar{\nu}_e) - F(\bar{\nu}_x)}$$

Flavor asymmetry

[Esteban-Pretel et al., 0706.2498]



← decoherence

● Flavor equilibration in both NH & IH

Quasi single-angle behaviour

● Nothing occurs in NH (stable configuration)

● Complete conversions in IH (bimodal instability)

Large  $\nu_e \bar{\nu}_e$  asymmetry required to suppress multi-angle decoherence



# MULTI-AZIMUTHAL-ANGLE (MAA) INSTABILITY

- Self-induced flavor conversions are associated to an instability in the flavor space [*Sawyer, 0803.4319; Banerjee, Dighe & Raffelt, 1107.2308*]
- Instability required to get started (exponential growth of the off-diagonal density matrix part)
- The onset of the conversions can be found through a stability analysis of the linearized EOMs.


In [*Raffelt, Sarikas, Seixas, 1305.7140*] a stability analysis of the EOMs has been performed including the azimuthal angle  $\phi$  of the  $\nu$  propagation and without enforcing axial symmetry.

A new multi-azimuthal-angle (MAA) instability has been found (also starting with a completely symmetric  $\nu$  emission) !!

- In the unstable case, numerical simulations are mandatory.

# MAA EFFECTS FOR SN NEUTRINOS

[A.M., 1308.1402]



Monochromatic  $\nu$  beam

System of only  $\nu_e$  and  $\bar{\nu}_e$  with an excess of  $\nu_e$

— MZA & MAA


- - - SZA & MAA

NH becomes unstable under MAA effects!


SZA vs MZA marginal impact

New effects in IH only for small  $\varepsilon$

# TOY MODEL NEUTRINO SPECTRA




System of only  $\nu_e$  and  $\bar{\nu}_e$  with an excess of  $\nu_e$



Fermi-Dirac energy distributions with  $\langle E \rangle = 15$  MeV


$$g_\omega \equiv \frac{|\Delta m^2|}{2\omega^2} \times \begin{cases} f_{\nu_e}(E_\omega) - f_{\nu_\mu}(E_\omega) & \text{for } \omega > 0 \\ f_{\bar{\nu}_\mu}(E_\omega) - f_{\bar{\nu}_e}(E_\omega) & \text{for } \omega < 0 \end{cases}$$



$$E_\omega = \left| \frac{\Delta m^2}{2\omega} \right|$$

# SPECTRAL SPLITS ( $\varepsilon=1.5$ )


[A.M., 1308.1402]



- Spectral swaps and splits in both NH (MAA) & IH (Bimodal)
- Antinu split smeared by MZA effects

# SPECTRAL SPLITS ( $\varepsilon=0.5$ )


[A.M., 1308.1402]



- Spectral swaps and splits IH (lead by the bimodal instability)
- Spectral swaps and splits NH **washed out**

# SPECTRAL SPLITS ( $\varepsilon=0.3$ )


[A.M., 1308.1402]



- Spectral swaps and splits IH **strongly smeared** (by MAA effects)
- Spectral swaps and splits IH **washed out**

# SPECTRAL SPLITS FOR SN NEUTRINO FLUXES

[A.M., 1308.5255]



(accretion phase)

$$F_{\nu_e} : F_{\bar{\nu}_e} : F_{\nu_x} = 2.4 : 1.6 : 1.0$$

$$\langle E_e \rangle = 12 \text{ MeV};$$

$$\langle E_{\bar{e}} \rangle = 15 \text{ MeV};$$


$$\langle E_x \rangle = 18 \text{ MeV}$$

$$\varepsilon = 1.34$$

- Spectral swaps and splits in both NH & IH !!
- In the axial symmetric case, only IH unstable

# SPECTRAL SPLITS FOR SN NEUTRINO FLUXES

[A.M., 1308.5255]



- Nu fluxes present a spectral split at  $E = 12$  MeV (fixed by lepton number conservation)
- Antinu fluxes are swapped



# CONCLUSIONS

- A new multi-azimuthal-angle (MAA) instability emerges in the dense SN neutrino gas, removing the axial symmetry in the nu-nu interaction term
- For simple systems of only  $\nu_e$  and  $\bar{\nu}_e$ , with an excess of  $\nu_e$ , MAA effects can lead to spectral splits (for large flavor asymmetry) or flavor decoherence (for small flavor asymmetry) in NH.
- For SN  $\nu$  spectra as in the accretion phase, MAA effects would induce swaps and splits in NH.
- A change of paradigm in the SN neutrino self-induced effects?

Still many open points (e.g. role of dense ordinary matter, initial nu angular distributions...)

LOT OF WORK TO BE DONE ...WAITING THE NEXT SN EXPLOSION !